

Hamiltonians and Green's functions which interpolate between two and three dimensions

Rainer Dick¹

*Department of Physics and Engineering Physics, University of Saskatchewan,
116 Science Place, Saskatoon, SK S7N 5E2, Canada*

Abstract

I propose to use Hamiltonians which contain two-dimensional and three-dimensional terms for the description of two-dimensional systems in physics.

As a model system the evolution of three-dimensional wavefunctions in the presence of an infinitely thin layer is studied. The model predicts distance laws for correlation functions which interpolate between two-dimensional and three-dimensional behavior. It also predicts that in certain cases transmission probabilities through thin layers should depend not only on the transverse, but also on the longitudinal momentum of the infalling particles.

The model also yields a static potential which interpolates between the two-dimensional logarithmic potential at small distances and the three-dimensional $1/r$ -potential at large distances.

1 Introduction

Two dimensions played a prominent role in the development of physics in the last twenty years.

On the experimental side this was driven e.g. by the needs of very large scale integration, by applications of semiconducting layer structures, by exploitations of surface catalytic effects, and by the development of atomic-scale surface analysis and manufacturing tools like scanning tunneling microscopy and atomic force microscopy, to mention only a few developments in this area.

On the theoretical side interest in two-dimensional field theories was largely driven by string theory [1], where fundamental excitations are described by covariant two-dimensional field theories, and by the realization that in two-dimensional critical systems with a rotational symmetry scaling symmetry may be elevated to full conformal invariance [2].

The purpose of the present paper is to point out that recent developments in the mathematical formulation of brane world models may also inspire new developments in

¹ rainer@sask.usask.ca

the physics of low-dimensional systems, and help us to acquire a better understanding of the transition between three-dimensional and two-dimensional behavior in these systems.

The present work was specifically motivated by the brane world model of Dvali, Gabadadze and Porrati, who recently proposed and analyzed a model which combined gravity on a $(3+1)$ -dimensional manifold (a "3-brane") with gravity in an ambient $(4+1)$ -dimensional bulk [3]. They observed that the combination of gravity in different dimensions yields a gravitational potential which interpolates continuously between the three-dimensional $-1/r$ potential at small distances and the four-dimensional $-1/r^2$ potential at large distances, with a transition scale $\ell_{DGP} \simeq m_3^2/m_4^3$ set by the ratios of the reduced Planck masses on the brane and in the bulk. Since we are not concerned with gravity in the present paper I will not explicitly write down the model in terms of intrinsic and extrinsic curvature terms, see [4], but instead refer to the related model of Dvali, Gabadadze and Shifman [5], which combines a Maxwell term in Minkowski space (coordinates $x^\mu = \{t, \mathbf{r}\}$) with a Maxwell term in an ambient $(4+1)$ -dimensional bulk (coordinates $x^M = \{t, \mathbf{r}, x^\perp\}$):

$$S = - \frac{1}{4q_3^2} \int dt \int d^3\mathbf{r} F_{\mu\nu} F^{\mu\nu} \Big|_{x^\perp=0} - \frac{1}{4q_4^2} \int dt \int d^3\mathbf{r} \int dx^\perp F_{MN} F^{MN}. \quad (1)$$

The resulting Coulomb potential on the $(3+1)$ -dimensional Minkowski space² is [5]

$$A^0(\mathbf{r}) = \frac{q_3}{4\pi r} \left[\cos\left(\frac{2q_3^2}{q_4^2} r\right) - \frac{2}{\pi} \cos\left(\frac{2q_3^2}{q_4^2} r\right) \text{Si}\left(\frac{2q_3^2}{q_4^2} r\right) + \frac{2}{\pi} \sin\left(\frac{2q_3^2}{q_4^2} r\right) \text{ci}\left(\frac{2q_3^2}{q_4^2} r\right) \right], \quad (2)$$

with the sine and cosine integrals

$$\text{Si}(x) = \int_0^x d\xi \frac{\sin \xi}{\xi}, \quad \text{ci}(x) = - \int_x^\infty d\xi \frac{\cos \xi}{\xi}.$$

The ratio of gauge couplings defines a length scale

$$\ell = \frac{q_4^2}{2q_3^2}, \quad (3)$$

and A^0 interpolates between a three-dimensional distance law at short distances and a four-dimensional distance law at large distances:

$$\begin{aligned} r \ll \ell : \quad A^0(\mathbf{r}) &= \frac{q_3}{4\pi r} \left[1 + \frac{2r}{\pi\ell} \left(\gamma - 1 + \ln\left(\frac{r}{\ell}\right) \right) + \mathcal{O}\left(\frac{r^2}{\ell^2}\right) \right], \\ r \gg \ell : \quad A^0(\mathbf{r}) &= \frac{q_3\ell}{2\pi^2 r^2} \left[1 - 2\frac{\ell^2}{r^2} + \mathcal{O}\left(\frac{\ell^4}{r^4}\right) \right]. \end{aligned}$$

Action principles of the kind (1) were denoted as dimensionally hybrid action principles in [6].

² See Ref. [6] for a discussion of dynamical potentials on the brane and in the bulk.

Of course, this does not simply carry over to low-dimensional systems in condensed matter or statistical physics: Dimensionally hybrid action principles would not be a suitable tool for model building in theoretical investigations of these systems because time derivatives generically will appear only as bulk terms in the Lagrangian of a particle interacting with a low-dimensional structure.

Therefore the main proposal of the present work is to use *dimensionally hybrid Hamiltonians* in model building for low-dimensional systems: A combination of two-dimensional and three-dimensional terms and the ensuing interpolating correlation functions may help to narrow the gap between the powerful methods of two-dimensional field theory and realistic thin layers or surface structures in physics and technology.

In the sequel I will use this idea to discuss non-relativistic particles interacting with a thin layer. The system becomes a dimensionally hybrid system with a specifically two-dimensional component through the assumption that particles in the layer have a kinetic energy different from particles outside of the layer, e.g. as a consequence of mass renormalization $M \rightarrow m$ due to the interaction of the particles with the components of the layer. This yields some straightforward but interesting results.

In the next section I will show in a simple model that transmission probabilities through thin layers in this class of models depend also on the momentum parallel to the layer if $\mu = m(L)/L|_{L \rightarrow 0}$ remains finite. The discussion of Green's functions in these models will be the subject of Secs. 3 and 4. Sec. 3 contains in particular a potential which interpolates between two-dimensional and three-dimensional distance laws.

2 Dimensionally hybrid Hamiltonians and Green's functions

To investigate implications of dimensionally hybrid Hamiltonians for the description of the interaction of particles with a thin layer we assume in the present section that the layer is planar and homogeneous and therefore generates a potential $U(z)$, where z is transverse to the layer.

In realistic two-dimensional systems particles are not strictly bound to the layer, and the effective particle mass in the layer may be changed due to interactions. This motivates the following Hamiltonian for particles of mass M in the presence of the layer

$$H = \frac{\hbar^2}{2\mu} \int d^2\mathbf{x} \nabla\psi^+ \cdot \nabla\psi \Big|_{z=0} + \int d^2\mathbf{x} \int dz \left(\frac{\hbar^2}{2M} (\nabla\psi^+ \cdot \nabla\psi + \partial_z\psi^+ \cdot \partial_z\psi) + \psi^+ U\psi \right). \quad (4)$$

Here and in the sequel all vectors are 2-dimensional vectors in the layer.

The assumption behind the Hamiltonian (4) is that the same field ψ may describe e.g. free electrons in the bulk and large polarons³ or other collective excitations involving conduction electrons in the layer. The parameter μ has dimensions of mass per length, and in a limiting procedure from layers of finite thickness L would correspond to

³ See e.g. [7,8,9] for introductions to polarons in solids.

$$\mu = \lim_{L \rightarrow 0} \frac{m(L)}{L}, \quad (5)$$

where $m(L)$ would be the mass of the modes in the layer.

The corresponding equation of motion for stationary single-(quasi)particle wavefunctions is

$$E\psi(\mathbf{x}, z) = -\delta(z)\frac{\hbar^2}{2\mu}\Delta\psi(\mathbf{x}, 0) - \frac{\hbar^2}{2M}\left(\Delta + \partial_z^2\right)\psi(\mathbf{x}, z) + U(z)\psi(\mathbf{x}, z). \quad (6)$$

The Fourier *ansatz*

$$\psi(\mathbf{x}, z) = \frac{1}{2\pi} \int d^2\mathbf{k} \psi(\mathbf{k}, z) \exp(i\mathbf{k} \cdot \mathbf{x}) \quad (7)$$

yields the separated equation

$$\left(E - \frac{\hbar^2 k^2}{2M}\right)\psi(\mathbf{k}, z) = -\frac{\hbar^2}{2M}\partial_z^2\psi(\mathbf{k}, z) + U(z)\psi(\mathbf{k}, z) + \delta(z)\frac{\hbar^2 k^2}{2\mu}\psi(\mathbf{k}, 0). \quad (8)$$

Every solvable model of one-dimensional quantum mechanics gives a solution to this class of layer models, with the kinetic term of the layer modes only generating a cusp proportional to $(M/\mu)k^2$ in $\ln \psi(\mathbf{k}, z)$.

Obviously, the large longitudinal momentum modes are strongly affected by the existence of layer modes, but we will see in a moment that the two-dimensional kinetic term can also have a strong impact on modes with small longitudinal momentum.




Fig. 1: Contribution of a virtual planar mode to a particle penetrating a thin homogeneous layer.

It is a trivial exercise to adapt solutions of one-dimensional quantum mechanics to the cusp imposed by the layer modes, but it may be worthwhile to write down the modifications of the transmission coefficient due to the layer modes when the layer potential represents a work function: $U(z) = -w\delta(z)$. The transmission coefficient for an infalling particle of momentum $\{\mathbf{k}, k_\perp\}$ is

$$T(\mathbf{k}, k_\perp) = \left[1 + \frac{M^2}{k_\perp^2} \left(\frac{w}{\hbar^2} - \frac{k^2}{2\mu}\right)^2\right]^{-1}, \quad (9)$$

i.e. the layer modes increase the transmission probability for low longitudinal momentum modes $0 < \hbar^2 k^2 < 4\mu w$, and decrease the transmission probabilities for the modes of higher longitudinal momentum.

This model also predicts a resonance in transmission for a certain value $\hbar^2 k^2 = 2\mu w$ of the *longitudinal* momentum. This is as a genuine consequence of the two-dimensional kinetic term in (4) and may be the simplest way to test the viability of the idea of dimensionally hybrid Hamiltonians in low-dimensional systems.

Obviously, the prediction of dependence of transmission probabilities on longitudinal momenta requires finiteness of the parameter μ (5), i.e. a derivation of the "phenomenological" Hamiltonian (4) from a limiting procedure of purely three-dimensional models will require a thorough study of finite size effects on mass renormalization in solids.

3 Correlations on a layer

A model similar to (4) allows for a neat discussion of the impact of combinations of kinetic terms from different dimensions on the "two-dimensional" correlation functions on the layer.

For this we assume that the layer is not necessarily homogeneous, but generates a strongly localized potential

$$V(\mathbf{x}, z) = u(\mathbf{x})\delta(z)$$

along the layer:

$$H = \frac{\hbar^2}{2M} \int d^2\mathbf{x} \int dz \left(\nabla\psi^+ \cdot \nabla\psi + \partial_z\psi^+ \cdot \partial_z\psi \right) + \int d^2\mathbf{x} \left(\frac{\hbar^2}{2\mu} \nabla\psi^+ \cdot \nabla\psi + \psi^+ u\psi \right) \Big|_{z=0}. \quad (10)$$

The generating functional for correlation functions on the layer is⁴

$$\begin{aligned} Z[j, j^+] &= \int d\psi d\psi^+ \exp\left(-\beta H[\psi, \psi^+] - \int d^2\mathbf{x} [\psi^+(\mathbf{x}, 0)j(\mathbf{x}) + j^+(\mathbf{x})\psi(\mathbf{x}, 0)]\right) \\ &= \exp\left(-\beta \int d^2\mathbf{x} \frac{\delta}{\delta j(\mathbf{x})} u(\mathbf{x}) \frac{\delta}{\delta j^+(\mathbf{x})}\right) Z_0[j, j^+] \end{aligned} \quad (11)$$

with

$$Z_0[j, j^+] \propto \exp\left(\frac{2M}{\hbar^2\beta} \int d^2\mathbf{x} \int d^2\mathbf{x}' j^+(\mathbf{x}) G(\mathbf{x} - \mathbf{x}', 0) j(\mathbf{x}')\right). \quad (12)$$

Since (10) is a free theory from a field theory point of view, the "two-dimensional" correlations in it can be calculated from tree-level diagrams, which involve only the restriction of the free Green's function $G(\mathbf{x} - \mathbf{x}', z)$ to the layer and insertions of the layer potential.

⁴ As usual $\delta/\delta j$ acts from the right if ψ is fermionic.

The free Green's function used in (12) satisfies

$$(\Delta + \partial_z^2) G(\mathbf{x}, z) + 2\ell\delta(z)\Delta G(\mathbf{x}, 0) = -\delta(\mathbf{x})\delta(z) \quad (13)$$

where

$$2\ell = \frac{M}{\mu}. \quad (14)$$

The *ansatz*

$$G(\mathbf{x}, z) = \frac{1}{(2\pi)^3} \int d^2\mathbf{k} \int dk_\perp G(\mathbf{k}, k_\perp) \exp[i(\mathbf{k} \cdot \mathbf{x} + k_\perp z)] \quad (15)$$

yields

$$(k^2 + k_\perp^2) G(\mathbf{k}, k_\perp) + \frac{\ell}{\pi} k^2 \int dk'_\perp G(\mathbf{k}, k'_\perp) = 1. \quad (16)$$

This determines the k_\perp -dependence of the propagator

$$G(\mathbf{k}, k_\perp) = \frac{f(k)}{k^2 + k_\perp^2}. \quad (17)$$

With

$$\frac{1}{\pi} \int_{-\infty}^{\infty} dk'_\perp \frac{1}{k^2 + k'^2_\perp} = \frac{1}{k}$$

we find the Green's function

$$G(\mathbf{k}, k_\perp) = \frac{1}{(1 + k\ell)(k^2 + k_\perp^2)} \quad (18)$$

or

$$G(\mathbf{k}, z) = \frac{1}{2k(1 + \ell k)} \exp(-k|z|). \quad (19)$$

The solution⁵

$$G(\mathbf{x}, z) = \frac{1}{8\pi^2} \int_0^\infty dk \int_0^{2\pi} d\varphi \frac{\exp[k(ir \cos \varphi - |z|)]}{1 + k\ell} = \frac{1}{4\pi} \int_0^\infty dk \frac{\exp(-k|z|)}{1 + k\ell} J_0(kr) \quad (20)$$

of (13) can be thought of as the electrostatic potential of a unit charge on the layer, if the fields which are continuous across the layer make a special contribution to the Hamiltonian

⁵ I follow the conventions of [10] for Bessel and Struve functions.

of the electromagnetic field⁶:

$$H[F] = \ell \int d^2\mathbf{x} \left(\mathbf{E}^2 + B_\perp^2 \right) + \frac{1}{2} \int d^2\mathbf{x} \int dz \left(\mathbf{E}^2 + E_\perp^2 + \mathbf{B}^2 + B_\perp^2 \right), \quad (21)$$

e.g. as a consequence of a non-vanishing limit of

$$2\ell = \lim_{L \rightarrow 0} (\epsilon_r L).$$

Here ϵ_r is the relative permittivity of the layer and L its transverse extension.

The perturbation series (11,12) requires the Green's function on the layer, which can be expressed as a linear combination of a Struve function and a Bessel function of the second kind:

$$\Phi(\mathbf{x}) = G(\mathbf{x}, z)|_{z=0} = \frac{1}{8\ell} \left[\mathbf{H}_0\left(\frac{r}{\ell}\right) - Y_0\left(\frac{r}{\ell}\right) \right]. \quad (22)$$


This interpolates between two-dimensional and three-dimensional distance laws

$$r \ll \ell : \quad \Phi(\mathbf{x}) = \frac{1}{4\pi\ell} \left[-\gamma - \ln\left(\frac{r}{2\ell}\right) + \frac{r}{\ell} + \mathcal{O}\left(\frac{r^2}{\ell^2}\right) \right],$$

$$r \gg \ell : \quad \Phi(\mathbf{x}) = \frac{1}{4\pi r} \left[1 - \frac{\ell^2}{r^2} + \mathcal{O}\left(\frac{\ell^4}{r^4}\right) \right].$$

$\Phi(\mathbf{x})$ along with the limiting cases is plotted in Fig. 2.

Fig. 2: The solid line is the Green's function (22) on the layer as a function of $x = r/\ell$, in units of ℓ^{-1} . The upper dashed line is the three-dimensional $1/4\pi r$ potential, and the lower dashed line is the two-dimensional logarithmic potential.




⁶ In the spirit of the philosophy advocated in this paper the electromagnetic fields in (21) would simultaneously describe photons in the bulk and polaritons in the layer.

Polaron masses in semiconductors are of the order of effective band masses. Due to strong band curvature the ratios of polaron masses to the mass of free electrons can be of order $m/M \simeq 10^{-2}$, so one may hope that in thin semiconducting layers of thickness L the parameter ℓ is of order $\ell \simeq 10^2 L$. Two-dimensional distance laws for correlation functions might then be realized up to distances of order $10L$, and intermittent behavior for distances between $10L$ and $200L$. For a possible realization of (22) as an electromagnetic potential in thin layers, semiconducting compounds involving Pb might be the best bet due to their high relative permittivities of order $\epsilon_r \sim 10^2 - 10^3$ [11]. The corresponding field strength on the layer is

$$-\partial_r \Phi(\mathbf{x}) = \frac{1}{8\ell^2} \left[\mathbf{H}_1\left(\frac{r}{\ell}\right) - Y_1\left(\frac{r}{\ell}\right) - \frac{2}{\pi} \right]. \quad (23)$$

Fig. 3: The solid line is the field strength per charge (23) on the layer as a function of $x = r/\ell$, in units of ℓ^{-2} . The dashed line approaching the solid line for $x < 1$ is the two-dimensional $1/4\pi\ell r$ field, and the other dashed line is the three-dimensional $1/4\pi r^2$ field.



This reduction of the force between charges in a thin dielectric layer with finite $2\ell = \lim_{L \rightarrow 0}(\epsilon_r L)$ can pictorially be understood as a consequence of the fact that field lines are refracted away from the layer when they leave the layer. This reduces the field lines e.g. between two opposite charges at short distances, since the field lines cannot re-enter the layer on short scales, whereas for large separation of the two charges along the layer re-entry renders the refraction effect negligible.

4 The Green's function for scattering from two-dimensional potentials on the layer

The stationary wave equation from (10) is

$$E\psi(\mathbf{x}, z) = \delta(z) \left(-\frac{\hbar^2}{2\mu} \Delta + u(\mathbf{x}) \right) \psi(\mathbf{x}, 0) - \frac{\hbar^2}{2M} (\Delta + \partial_z^2) \psi(\mathbf{x}, z). \quad (24)$$

We have already noticed that this can be solved exactly for $u(\mathbf{x})=\text{constant}$.

In discussing scattering of bulk particles from the layer in the model (10) we could proceed using the ordinary three-dimensional Green's function for scattering of waves of energy $E = \hbar^2 K^2 / 2M$ and treat the full two-dimensional contribution to Eq. (24) as a perturbation. However, here I rather would like to treat only the layer potential $u(\mathbf{x})$ as a perturbation. This has the virtue of reducing the perturbation for large longitudinal momenta.

The relevant unperturbed wave is then

$$\begin{aligned} \psi_0(\mathbf{x}, z) = \frac{1}{\sqrt{2\pi^3}} \exp(i\mathbf{K}_{\parallel} \cdot \mathbf{x}) & \left[\Theta(-z) \left(\exp(iK_{\perp} z) + \frac{K_{\parallel}^2 \ell}{iK_{\perp} - K_{\parallel}^2 \ell} \exp(-iK_{\perp} z) \right) \right. \\ & \left. + \Theta(z) \frac{K_{\perp}}{K_{\perp} + iK_{\parallel}^2 \ell} \exp(iK_{\perp} z) \right] \end{aligned} \quad (25)$$

where again the definition (14) was used.

The relevant Green's function G_K for propagation of bulk plane waves of energy

$$E = \frac{\hbar^2 K^2}{2M} = \frac{\hbar^2}{2M} (\mathbf{K}_{\parallel}^2 + K_{\perp}^2)$$

has to satisfy

$$(\Delta + \partial_z^2 + K^2) G_K(\mathbf{x}, z) + 2\ell\delta(z)\Delta G_K(\mathbf{x}, 0) = -\delta(\mathbf{x})\delta(z), \quad (26)$$

and the solution proceeds similarly to the solution of (13). The Fourier *ansatz* (15) yields

$$(k^2 + k_{\perp}^2 - K^2) G_K(\mathbf{k}, k_{\perp}) + \frac{\ell}{\pi} k^2 \int dk'_{\perp} G_K(\mathbf{k}, k'_{\perp}) = 1, \quad (27)$$

which determines the k_{\perp} -dependence of the propagator

$$G_K(\mathbf{k}, k_{\perp}) = \frac{f(k)}{k^2 + k_{\perp}^2 - K^2}. \quad (28)$$

At this stage the possibility of poles complicates the calculation slightly: In evaluating the integral in (27) with (28) for $k < K$ we have to make a judicious choice on how to shift the poles or the integration path at $k_{\perp} = \pm\sqrt{K^2 - k^2}$, corresponding to correct physical boundary conditions on $G_K(\mathbf{k}, z)$. The correct choice turns out to be $k_{\perp} = \pm(\sqrt{K^2 - k^2} + i\epsilon)$ since $G_K(\mathbf{k}, z)$ is supposed to describe outgoing scattered waves from the layer if $k < K$, i.e. we have

$$\begin{aligned} G_K(\mathbf{k}, k_{\perp}) &= \frac{f(k)}{k^2 + k_{\perp}^2 - K^2 - i\epsilon} \\ &= f(k) \left(\mathcal{P} \frac{1}{k^2 + k_{\perp}^2 - K^2} + i\pi\delta(k^2 + k_{\perp}^2 - K^2) \right). \end{aligned} \quad (29)$$

With

$$\frac{1}{\pi} \int_{-\infty}^{\infty} dk'_{\perp} \frac{1}{k^2 + k'^2_{\perp} - K^2 - i\epsilon} = \frac{\Theta(k^2 - K^2)}{\sqrt{k^2 - K^2}} + i \frac{\Theta(K^2 - k^2)}{\sqrt{K^2 - k^2}}$$

we find the Green's function at large longitudinal wavelength $k < K$

$$G_K(\mathbf{k}, k_{\perp}) = \frac{\sqrt{K^2 - k^2}}{(\sqrt{K^2 - k^2} + ik^2\ell)(k^2 + k_{\perp}^2 - K^2 - i\epsilon)}, \quad (30)$$

$$G_K(\mathbf{k}, z) = \frac{1}{2k^2\ell - 2i\sqrt{K^2 - k^2}} \exp(i\sqrt{K^2 - k^2}|z|), \quad (31)$$

while the short longitudinal wavelength part is

$$G_K(\mathbf{k}, k_{\perp}) = \frac{\sqrt{k^2 - K^2}}{(\sqrt{k^2 - K^2} + k^2\ell)(k^2 + k_{\perp}^2 - K^2 - i\epsilon)}, \quad (32)$$

$$G_K(\mathbf{k}, z) = \frac{1}{2\sqrt{k^2 - K^2} + 2k^2\ell} \exp(-\sqrt{k^2 - K^2}|z|). \quad (33)$$

Of course, the Green's function again reduces to the usual three-dimensional result for $\ell \rightarrow 0$, i.e. if the modes in the layer become so heavy that they decouple.

With an incoming plane wave the integral equation following from (24,26) is

$$\psi(\mathbf{x}, z) = \psi_0(\mathbf{x}, z) - \frac{2M}{\hbar^2} \int d^2\mathbf{x}' G_K(\mathbf{x} - \mathbf{x}', z) u(\mathbf{x}') \psi(\mathbf{x}', 0). \quad (34)$$

In a Born approximation this yields with (25)

$$\psi(\mathbf{x}, z) = \psi_0(\mathbf{x}, z) - \frac{2M}{\sqrt{2\pi}^7 \hbar^2} \frac{K_{\perp}}{K_{\perp} + iK_{\parallel}^2\ell} \int d^2\mathbf{k} \exp(i\mathbf{k} \cdot \mathbf{x}) G_K(\mathbf{k}, z) u(\mathbf{k} - \mathbf{K}_{\parallel}), \quad (35)$$

where the normalization of the Fourier transformed layer potential is

$$u(\mathbf{q}) = \int d^2\mathbf{x} \exp(-i\mathbf{q} \cdot \mathbf{x}) u(\mathbf{x}).$$

Eq. (35) together with (33) implies that no particular scattering wave is generated by the short wavelength components at $|\mathbf{q}| < K_{\parallel} + \sqrt{K_{\parallel}^2 + K_{\perp}^2}$ of the layer potential, which is simply a statement of the limited resolving power of the external wave. However, beyond that the ℓ -dependence of (31) and (35) implies that the two-dimensional kinetic term in (10) reduces potential scattering at large K_{\parallel} .

5 Summary

Two-dimensional field theory is a very appealing subject with many powerful results. One reason for this is because every tensor and spinor field on a 2-manifold decomposes

into covariant primary fields which provide 1-dimensional representations of the corresponding symmetry groups⁷.

However, the assumption of strict two-dimensionality seems too restrictive when it comes to comparisons with actual layers or surface structures in physics. Conservative theoretical approaches to low-dimensional structures in physics and technology therefore rely on genuine three-dimensional Hamiltonians and only restrict the locations and momenta of particles to a surface or a layer (see e.g. Sec. 9.2 in [8]). In these approaches two-dimensionality is only taken into account at a kinematical level, at the expense of sacrificing the powerful methods and results of two-dimensional field theory.

On the other hand, recent results in brane theory taught us that straightforward combinations of four-dimensional terms and five-dimensional terms in action principles yield interpolating Green's functions on the brane, and it is apparent from the functional integral representation that this property must also hold for higher order correlation functions on the brane.

As mentioned above this cannot carry over directly to low-dimensional systems in condensed matter and statistical physics, but it initiated the present proposal to use dimensionally hybrid Hamiltonians $H = \ell h_2 + H_3$ for theoretical investigations of low-dimensional structures in physics.

Such an approach has the prospect to provide more realistic results than strictly two-dimensional field theory, while at the same time utilizing the power of two-dimensional field theory for the determination e.g. of equilibrium correlation functions in the limiting cases $\ell \rightarrow \infty$ or $k\ell \gg 1$.

Further virtues of this approach are predictions on the transition behavior between two-dimensional and three-dimensional distance laws in layers and surface structures, and a better understanding on how two-dimensional structures might be approached in the more conservative purely three-dimensional framework, through studies of finite size effects on effective masses and permittivities in three-dimensional models.

To illustrate the use and some straightforward consequences of dimensionally hybrid Hamiltonians a homogeneous layer and layers with strongly localized potentials were studied. In these settings the field ψ describes simultaneously bulk particles of mass M and excitations of mass μ per transverse length in the layers. These models might serve as an approximation to thin layers of semiconductors or polar solids, where the planar modes would correspond to large polarons. The model predicts a strong dependence of transmission probabilities on longitudinal momentum.

The Green's function in the model also yields a static potential which interpolates between the logarithmic two-dimensional distance law at distances $\ll \ell$ and the three-dimensional Coulomb law at distances $\gg \ell$.

Acknowledgement: This work was supported by NSERC.

⁷ This is true beyond the realm of conformal transformations if the Beltrami parameters on the 2-manifold are used to decompose tensors and spinors into *covariant primary fields*, see Secs. 1 and 2 in [12] for tensors and [13] for spinors.

References

- [1] M.B. Green, J.H. Schwarz, E. Witten, *Superstring Theory* (Cambridge University Press, Cambridge, 1987).
- [2] A.A. Belavin, A.M. Polyakov, A.B. Zamolodchikov, *Infinite conformal symmetry in two-dimensional quantum field theory*, Nucl. Phys. B 241 (1984) 333–380.
- [3] G. Dvali, G. Gabadadze, M. Porrati, *4-D gravity on a brane in 5-D Minkowski space*, Phys. Lett. B 485 (2000) 208–214.
- [4] R. Dick, *Brane worlds*, Class. Quantum Grav. 18 (2001) R1–R23.
- [5] G. Dvali, G. Gabadadze, M. Shifman, *(Quasi)localized gauge field on brane: dissipating cosmic radiation to extra dimensions?*, Phys. Lett. B 497 (2001) 271–280.
- [6] R. Dick, D.M.E. McArthur, *A photon mass on the brane*, hep-th/0203271, Phys. Lett. B (in press).
- [7] C. Kittel, *Quantum Theory of Solids*, 2nd ed. (Wiley, New York, 1987).
- [8] O. Madelung, *Introduction to Solid-State Theory*, 3rd pr. (Springer, Berlin, 1996).
- [9] J.T. Devreese, *Polarons*, in: *Encyclopedia of Applied Physics*, Vol. 14, eds. G.L. Trigg and E.H. Immergut (Wiley-VCH, New York, 1995) pp. 383–409.
- [10] M. Abramowitz, I.A. Stegun, *Handbook of Mathematical Functions* (United States National Bureau of Standards, Washington, 1972).
- [11] Landolt–Börnstein *Numerical Data and Functional Relationships in Science and Technology* Vol. III/17f, ed. O. Madelung (Springer, Berlin, 1983).
- [12] R. Dick, *Topological aspects of chiral fields in two dimensions and superstring vertices*, Fortschr. Phys. 40 (1992) 519–591.
- [13] H. Nicolai, *New linear systems for 2D Poincaré supergravities*, Nucl. Phys. B 414 (1994) 299–328.